BULGARIAN SOCIETY OF RHEUMATOLOGY

CONSENSUS

PROGRESSIVE SYSTEMIC SCLEROSIS

Prof. Dr. R. Rashkov, MD

Dr. V. Reshkova, MD

Dr. D. Kalinova, MD

Dr. V. Boyadzhieva, MD

Dr. Tsv. Yoneva

March, 2019

DEFINITION

Systemic sclerosis is a clinically heterogeneous, multisystemic, progressive autoimmune disease characterized by impaired microcirculation and connective tissue deposition in the skin and internal organs.

Systemic sclerosis is characterized by a variety of clinical symptoms and syndromes described in Table 1.

Table 1. Pathological manifestations of systemic sclerosis (Y. Shoenfeld et al., 2012)

Skin	Raynaud's syndrome; sclerodactyly; ulcers on the fingertips; scleroderma of the limbs, face, neck, torso; telangiectasia; calcification
Lungs	Basal pulmonary fibrosis; pulmonary hypertension
Heart	Arrhythmias; myocardial fibrosis; congestive heart failure
Musculoskeletal system	Arthritis/arthritis, Tendon friction ribs; joint contractures; myopathy
Gastrointestinal tract	Esophagopathy; small intestinal hypomotility
Kidneys	Scleroderma renal crisis (malignant hypertension, rapidly progressive renal failure)

CLASSIFICATION

Skin involvement, also called scleroderma, is the most characteristic clinical symptom of systemic sclerosis. Scleroderma is divided into two main forms – localized scleroderma, limited to the skin and subcutaneous tissue, and systemic sclerosis,

affecting the skin and internal organs. In turn, systemic sclerosis is divided into two types - limited and diffuse. Limited systemic sclerosis is manifested by scleroderma mainly on the hands, forearms, less on the face and neck. Diffuse systemic sclerosis is characterized by diffuse skin sclerosis and involvement of various internal organs.

Localized forms of scleroderma are characterized by limited sclerosis of the skin, which are distinguished by the morphology of dermal lesions. There are two main types: linear scleroderma and morphea. Linear scleroderma is characterized by the presence of longitudinal plaques of cutaneous sclerosis, while in morphea limited oval plaques are observed. Specific forms of linear scleroderma occur when the head and face are involved, called "en coup de saber", which cause severe facial and cranial deformities, as well as neurological abnormalities. Several basic forms of morphea have been described: limited, which is divided into superficial and deep subtypes, generalized morphea, and pansclerotic morphea. Some rare forms that belong to the spectrum that covers the morphea have also been described: keloid or nodular morphea (sclerotic papules or plaques resembling keloid scars), Pasini and Pierini atrophoderma (hyperpigmented plaques, with mild skin depression, demarcated border and lack of inflammation and compaction) and lichen sclerosus et atrophicus (atrophic white or shiny plaques that may indicate follicular obstruction).

Fig. 1.

*Morphea – poster morphea, diffuse morphea, morphea profunda, pansclerotic morphea, Pasini/Pierini atrophoderma, nodular morphea

**CREST – calcinosis, Raynaud's syndrome, esophagitis, sclerodactyly, telangiectasia

Linear scleroderma is more common in children, while morphea is more common in the adult population.

The expected annual incidence is 2.7 per 100,000, with a prevalence of 50 per 100,000; The onset of the disease is around the age of 46, and the ratio of women to men is 2.6. In children with localized scleroderma, treatment is often delayed due to a misdiagnosis.

MORPHEA

The limited form of morphea, also known as morphea en plaque, is the most common form of morphea, accounting for about 50% of all reported cases. Lesions usually consist of slowly growing single or numerous limited plaques on the body or limbs. The described morphea can be divided into superficial and deep subtypes, with superficial being the most common. In the superficial variant, plaque-type lesions are characterized as well-defined, compacted plaques that range from 1 to more than 20 centimetres in diameter. These lesions are relatively superficial, involving mainly the dermis. They often start as oval, erythematous or slightly edematous plaques, which when the disease is active have a purple colour on the edge of the lesion and usually become more hyperpigmented over time. The involved skin looks shiny and there is a lack of sweat glands and hair follicles in the affected area. The histological result of an "early" lesion shows a dense inflammatory infiltration com-

posed of lymphocytes, macrophages, plasma cells, eosinophils and mast cells. The later stages are characterized by mild inflammation and deposition of collagen (type I and type III), extending from the reticular dermis to more superficial structures – indistinguishable from the biopsy finding in patients with systemic sclerosis.

Morphea profound subtype (morphea profunda)

The deep subtype of the limited form of morphea (formerly called subcutaneous morphea or morphea profunda) includes several clinical units in which inflammation and sclerosis occur in the deep dermis, panicle, fascia, and even the superficial muscle. It is characterized by limited, sclerotic plaques with the appearance of "pseudo-cellulite". The lesions are often hyperpigmented, but due to the deeper level of inflammation, they do not have other colour changes typical of the superficial subtype. As a complication of pansclerotic morphea, chronic ulcers have been observed that may evolve into squamous cell carcinoma.

Generalized morphea

The generalized form of morphea is characterized by the involvement of a large part of the skin surface with numerous induced plaques with hyperpigmentation and frequent muscle atrophy. It represents approximately 15% of all reported cases of morphea.

Diagnostic criteria:

The most commonly used diagnostic criteria are those of Takehara and Sato:

four or more plaques larger than 3 cm in diameter, whether of the type of oval or linear lesions

engage two or more than seven anatomical areas of the body, the seven areas are: head and neck, right upper limb, left upper limb, front of the torso, back of the trunk, right lower limb

The diagnosis is accepted in the presence of criterion 1) and/or criterion 2).

Numerous lesions of the generalized form often range from hyperpigmented to silvery in colour. Unlike systemic scleroderma, patients do not have sclerodactyly, Raynaud's phenomenon, capillaroscopic changes. The presence of a variety of antibodies: ANA, anti-Histones, anti-Nucleosomes, U3-RNP, fibrillin, anti-topoisomerase-II, ACL suggests that morphea is an autoimmune disease. In addition, the cytokine profile of the lesions includes activation of TH2 (T-helper type 2) and overexpression of pro-fibrotic cytokines, such as transforming growth factor β (TGF- β) and connective tissue-derived growth factor, similar to the profile observed in systemic sclerosis. Connective tissue growth factor is a soluble mediator that enhances and prolongs the profibrotic effects of TGF- β . The end result of endothelial damage and the inflammatory cascade is increased deposition of collagen and extracellular matrix. Other proposed pathophysiological mechanisms in morphea include the formation of antimatrix metalloproteinase antibodies, as well as increased expression of insulin-like growth factor, which improves collagen production.

Linear scleroderma

Linear scleroderma is considered in some classifications as a form of morphea (linear morphea), but given its clinical and morphological characteristics, it is accepted to be considered as a separate nosological unit. It is most commonly diagnosed in children and adolescents, representing most cases of localized scleroderma. Linear scleroderma is characterized by hypo- or hyperpigmented, compacted linear plaques, which are most often single, unilateral in 75-95% of cases, usually in the limbs. Older lesions may be atrophic or sclerotic. The changes cover the deep dermis, subcutaneous fat, muscles and bones and cause significant deformities. The involvement of the face and scalp from linear scleroderma is called "en coup de saber" and is associated with Parry-Romberg syndrome (hemifacial atrophy of the tissues under the forehead with

little involvement of the skin). "En coup de saber" is characterized by a linear, atrophic depression affecting the frontoparietal aspect of the face and scalp. With the participation of the scalp, alopecia is found, and in this variant, loss of eyebrows and eyelashes can be observed.

Progressive hemifacial atrophy, also known as Parry-Romberg syndrome, is a severe, segmental form of craniofacial linear scleroderma. Unlike "en coup de saber", the primary abnormality is found in subcutaneous fat, muscle and bone. Although the skin is usually not strengthened or bound, approximately 71% of patients develop skin sclerosis resembling "en coup de saber".

Both en coup de saber and Parry-Romberg syndrome can be associated with abnormalities in the CNS, ocular and dental status. Epileptic seizures occur in 13% of patients with both syndromes, according to some authors.

Table 2. Comparison between localized scleroderma and progressive systemic sclerosis (M. Mayes et al., 2011)

Clinical manifestations	Localized scleroderma	Progressive systemic sclerosis
Skin manifestations	Plaques or linear skin seal	Sclerodactyly $\pm$ proximal scleroderma
Raynaud's syndrome	–	+
Digital ischemic changes	–	+
Involvement of internal organs	–	+
Antinuclear antibodies	(+) $\geq$ 50% from the sick	(+) $\geq$ 90% from the sick
Scleroderma-specific antibodies	–	(+) around 60% from the sick
Histological finding	Deposition of collagen deposits in the dermis	Deposition of collagen deposits in the dermis

The approach to systemic sclerosis (SSc) has changed over the years with a growing emphasis on very early diagnosis of the disease. All classifications distinguish limited cutaneous SSc from diffuse, SSc, and this division into subgroups is useful in clinical practice, but does not include very early symptoms in the initial development of the disease.

Due to the lack of sensitivity of the 1980 criteria and advances in SSc knowledge, the American College of Rheumatology (ACR) and the European League Against Rheumatism (EULAR) set up a committee to submit a joint proposal for new SSc classification criteria. The objectives are to develop criteria that 1) include a wider range of SSc, including patients who are in the early and late stages of

the disease process; 2) include vascular, immunological and fibrotic manifestations; 3) are applicable in everyday clinical practice and 4) are in accordance with the criteria used to diagnose SSc in clinical practice.

Despite the ACR/EULAR classification criteria for SSc adopted in 2013, for very early forms of the disease, the ones proposed by EUSTAR (European League Against Rheumatism Scleroderma Trial and Research) in 2011 remain relevant.

EUSTAR criteria (2011):

Criteria considered to be of high clinical relevance for the very early diagnosis of SSc:

1. Raynaud's phenomenon
2. Diffuse swollen fingers, with progression to sclerodactyly
3. Abnormal capillaroscopic finding with scleroderma picture
4. Positive anti-centromeric antibodies
5. Positive anti-topoisomerase-1 antibodies.

Criteria considered leading to early referral:

1. Raynaud's phenomenon
2. Puffy fingers
3. Positive ANA.

Preferred diagnostic tools for diagnosing a patient with very early SSc are the available specific antibodies – anti-centromeres and anti-topoisomerase-1 and capillaroscopy of the nail bed.

The reason for diagnosing very early SSc is based on the assumption that the use of these criteria will lead to earlier diagnosis and earlier treatment of SSc, which should improve the long-term outcomes of therapy in patients with SSc. The application of the criteria in everyday practice aims to diagnose very early SSc and reduce the economic burden of this disease, as well as to improve the quality of care, through earlier, easier and more reliable screening of persons at risk

ACR/EULAR classification criteria (2013) for systemic sclerosis (Table 3).

Scleroderma-associated antinuclear antibodies are also included in modern classification criteria.

Antibodies in systemic sclerosis: Antibodies that are positive in systemic sclerosis are divided into two groups. The first group included autoantibodies that were found to have a direct pathogenetic role because they were involved in the process of tissue damage in systemic sclerosis (Table 4).

The other group includes antibodies directed against various nuclear structures (Scl-70; CENT; Ku; RNA polymerase I; RNA polymerase III; U1RNP; nucleolar proteins – U3RNP, Th/To, PM/Scl) or this is the group of antinuclear antibodies. Approximately 96% of patients with systemic sclerosis have positive antinuclear antibodies (ANAs) tested by indirect immunofluorescence. In turn, scleroderma-associated

Table 3. ACR/EULAR (2013) classification criteria for systemic sclerosis

Criteria	Sub criteria	Score
Skin thickening (scleroderma) of the fingers, extending proximally from the metacarpophalangeal joints (sufficient criterion)	–	9
Finger skin seal (only the higher score is taken into account)	Puffy fingers Sclerodactyly (distal to metacarpophalangeal joints but proximal to proximal interphalangeal joints)	2 4
Lesions on the fingertips (only the higher score is taken into account)	Digital ulcerations on the fingertips Dotted scars on fingertips	2 3
Telangiectasia	–	2
Abnormal capillaroscopic picture	–	2
Pulmonary arterial hypertension and/or interstitial pulmonary fibrosis	Pulmonary arterial hypertension Interstitial lung disease	2 2
Raynaud's syndrome	–	3
Positive scleroderma-linked antinuclear antibodies (anticentromeric, antitopoisomerase (anti-Scl-70), anti-RNA polymerase III) (maximum score is 3)	Anti-center (anti-CENT) At Antitopoisomerase (anti-Scl-70) At Antibodies against RNA polymerase III	3 3 3

* These criteria are not applicable in patients in whom the skin seal has not affected the fingers, as well as in patients with scleroderma-like diseases (nephrogenic sclerosing fibrosis, generalized morphea, eosinophilic fasciitis, diabetic scleroderma, scleromyxedema, porphyria, erythromyrosis -versus-host disease).

* Patients with a total score ≥ 9 are diagnosed with systemic sclerosis.

antinuclear antibodies can be divided into: scleroderma-specific and scleroderma-associated antibodies.

Table 4. Autoantibodies involved in the pathogenetic process in systemic sclerosis (J. Varga et al., 2011)

Autoantibodies	Target antigens
Anti-endothelial At	Endothelial cells
Anti-fibroblast At	The membrane of fibroblasts
Anti-collagenase At	Matrix metalloproteinases
Anti-PDGFR (platelet-associated growth factor receptor)	PDGF cell surface receptor
Anti-fibrillin-1	Extracellular microfibrils

- Scleroderma-specific At – anti-Scl-70, anti-CENT, anti-RNA polymerase III, anti-U3RNP, anti-Th/To At

- Scleroderma-associated At – anti-PM/Scl, anti-Ku, anti-U1RNP, anti-Ro52, anti-Ro60 At

In 60-70% of patients with systemic sclerosis, one of the three most common antibodies in systemic sclerosis is positive – anti-centromeric (anti-CENT)

At, antitopoisomerase (anti-Scl-70) At or antibodies against RNA polymerase III, included as criterion in modern classification criteria (Table 5).

Scleroderma-specific antibodies have predictive value in terms of the degree and type of skin involvement (diffuse or limited), the type of organ damage and the outcome of the disease.

Additionally, antibodies in systemic sclerosis can be divided into two main groups – At, specific for the diffuse form of scleroderma and At, specific for the limited form (Figure 2), ie. in patients with scleroderma, in addition to clinical, an immunological classification can be made.

Scleroderma-associated At (anti-PM/Scl, anti-Ku and anti-U1RNP) At occurs mainly in patients with overlapping syndromes or overlapping rheumatic diseases, combining clinical manifestations and symptoms characterizing various systemic connective tissue diseases such as myositis, systemic sclerosis, systemic lupus erythematosus.

A link between scleroderma-associated antibodies and disease prognosis has been demonstrated (Fig. 3).

Table 5. Scleroderma-specific antibodies and their associated clinical manifestations (M. Mayes et al., 2011)

	anti-CENT	anti-Scl-70	anti-RNA polymerase III
Limited form of scleroderma	+++	++	++
Diffuse form of scleroderma	+	+++	+++
Interstitial lung disease	+	+++	++
Renal crisis	+	++	+++
Pulmonary arterial hypertension (PAH)	+++ prolonged course of the disease	no connection established	no connection established

Legend: rarely +; Less characteristic ++; characteristic +++

Fig. 2. Relationship between the type of skin involvement and the different antibodies in systemic sclerosis (K. Ho et al., Arthritis Res Ther, 2003)

Fig. 3. Prognosis and scleroderma-associated antibodies (K. Ho et al., Arthritis Res Ther, 2003)

The described autoantibodies, the established frequencies in systemic sclerosis and the corresponding clinical associations are summarized in Table 6.

The differential diagnosis of systemic sclerosis is broad and includes a wide variety of diseases and conditions that occur with chronic skin thickening (Table 7).

Table 6. Frequency of scleroderma-associated antibodies and relevant clinical associations (S. Nihtyanova et C. Denton, Nat. Rev. Rheumatol., 2010)

Antibodies Frequency (%) Clinical associations		
Anti-CENT At	16-39%	lcSSc; pulmonary arterial hypertension without pulmonary fibrosis; protective role for scleroderma renal crisis and pulmonary fibrosis
Anti-Scl-70 (anti-topo I) At	9-39%	dsSSc > lsSSc; pulmonary fibrosis; pronounced digital vasculopathy
Anti-RNA polymerase (anti-RNAP) At	4-25%	dsSSc; scleroderma renal crisis
Anti-Th/To At	1-7%	lcSSc; pulmonary fibrosis; pulmonary arterial hypertension
Anti-U3RNP At	1-6%	dsSSc > lsSSc; muscle involvement (myositis/myopathy) pulmonary arterial hypertension
Anti-PM/Scl At	0-6%	Overlapping syndrome – myositis; arthritis Pulmonary fibrosis
Anti-Ku At	1-3%	Muscle and joint engagement
Anti-U1RNP At	5-35%	Overlapping syndromes
Anti-U11/U12RNP At	1.6-5%	Pulmonary fibrosis

ISSc – limited form of systemic sclerosis; dsSSc - diuresis form of systemic sclerosis.

Table 7. Differential diagnosis of conditions with chronic skin thickening (R. Simms et al., 2011)

Systemic sclerosis	<ol style="list-style-type: none"> 1. Limited systemic sclerosis 2. Diffuse systemic sclerosis 3. Overlapping syndromes <ul style="list-style-type: none"> • Sharp's disease • Systemic sclerosis/Polymyositis - „scleromyositis“ • Systemic sclerosis/Rheumatoid arthritis • Systemic sclerosis/systemic lupus erythematosus
Localized scleroderma	<ol style="list-style-type: none"> 1. Morphea 2. Linear scleroderma
Scleroderma	<ol style="list-style-type: none"> 1. Buschke's sclerosis 2. Diabetic scleredema 3. Scleredema in newborns
Scleromyxedema	<ul style="list-style-type: none"> • Extracellular mucin deposits in the skin and internal organs • It is associated with the production of monoclonal IgG proteins
Eosinophilic syndromes	<ol style="list-style-type: none"> 1. Eosinophilic fasciitis (diffuse fasciitis with eosinophilia, Schulman's disease) 2. Eosinophilia-myalgia syndrome 3. Toxic oil syndrome
Metabolic diseases	<ol style="list-style-type: none"> 1. Porphyria cutanea tarda 2. Phenylketonuria 3. Werner's syndrome 4. Acromegaly
Pseudoscleroderma	<ul style="list-style-type: none"> • Local application of vitamin K, bleomycin, penicillin
Nephrogenic systemic sclerosis	<ul style="list-style-type: none"> • Progressive systemic fibrosis in patients with end-stage renal failure
Chronic graft-versus host disease (GVHD)	<ul style="list-style-type: none"> • Allogeneic hematopoietic stem cell transplantation
Rare diseases	<ol style="list-style-type: none"> 1. Pachydermoperiostitis 2. POEMS (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy)

Table 8. Distinctive features between scleroderma and some scleroderma-like conditions (R. Simms et al., 2011)

Disease	Skin changes and spread	Non-skin manifestations	Histopathological characteristic	Other manifestations
Localized scleroderma	Amorphous plaques (morphea), linear seal of the face, limbs, fingers are not affected	–	Mononuclear infiltrates, collagen deposits in the dermis	It is common in children, the linear form - „en coup de saber“ or Parry Romberg syndrome, often positive At
Scleroderma	Skin seal in the torso, back, neck	Rarely	Swollen collagen bundles, mucin deposits	–
Scleromyxedema	Seal, sclerodactyly, microstomy	Rarely	Excessive deposition of mucin deposits in the dermis	Elevated serum paraprotein (especially IgG-λ)
Nephrogenic systemic sclerosis	Limbs, fingers, symmetrical involvement	Common (muscles, fascia, lungs)	Proliferation of fibroblasts, mucin deposits	–
Eosinophilic fasciitis	Limbs, torso, fingers are not affected	Rarely, (muscle/myositis)	Collagen deposits in the dermis and hypodermis, the epidermis is not affected, possible inflammation in the muscles	Often peripheral eosinophilia

Evaluation of the disease activity of systemic sclerosis in daily rheumatological practice (Table 9).

Table 9. Assessment of disease activity

Organ localization	Methods used in everyday clinical practice to assess disease activity
Skin	Modified Rodnan skin score
Heart	Echocardiography Right cardiac catheterization 6-minute walking test (6MWT)
Lungs	Functional study of respiration with gas diffusion High resolution computed tomography
Kidneys	Creatinine clearance Blood pressure monitoring Serum creatinine
Digital ulcerations	Active search for ulcers on the fingertips
Musculoskeletal system	Study of serum creatine phosphokinase levels Number of painful joints
Biomarkers	Pro-inflammatory indicators – ESR, CRP, fibrinogen Serum BNP/NT-Pro BNP (N-terminal brain natriuretic peptide)

Screening for pulmonary hypertension (C. Denton, 2011)

Algorithm for staging and behavior in interstitial pulmonary fibrosis (C. Denton, 2011)

Legend: VK – vital capacity; FER – functional examination of respiration; HRCT – high resolution computed tomography

Algorithm for behaviour in scleroderma renal crisis (V. Steen, 2011)

Modified Rodnan skin score

The degree of skin thickening in 17 areas was assessed using a scale of 0-3, respectively:

- 0-normal skin
- 1-light thickening
- 2-moderate thickening
- 3-heavy thickening

Then add up the total number of points, the maximum number of points - 51 points. (17 areas x 3)

Treatment of progressive systemic sclerosis

Progressive systemic sclerosis is a systemic

connective tissue disease that is difficult to treat. Treatment of scleroderma is based on a definite diagnosis, determination of the subtype of progressive systemic sclerosis according to the immunological profile, stage of the disease, organ status, as well as the presence of overlapping clinical manifestations. Drugs for systemic sclerosis include:

- Immunomodulatory and antifibrotic therapy or these are disease-modifying drugs
- Medicines used in various organ lesions.

Algorithm for the treatment of progressive systemic sclerosis **MEDICATIONS FOR PROGRESSIVE SYSTEMIC SCLEROSIS****I. Immunomodulatory therapy****1. Cyclophosphamide (Endoxan®, Cytozar®)****1.1. Mechanism of action**

- leads to dose-dependent lymphopenia, which occurs through T- and B-lymphocyte depletion
- impairs the function of macrophages
- affects gene transcription, which leads to a decrease in the production of autoantibodies

1.2. The effectiveness of Cyclophosphamide has been proven in:

- SSc-associated pulmonary fibrosis
- Digital ulcerations

2. Mycophenolate mofetyl (MMF), (Cellcept®)**2.1. Mechanism of action**

– inhibits inosine-5'-monophosphate dehydrogenase, inhibits the proliferation of T- and B-lymphocytes, respectively cell-mediated and humoral immune response

2.2. Proven efficacy on cutaneous (scleroderma) and pulmonary (fibrosis) manifestations in patients with progressive systemic sclerosis.

3. Methotrexate**3.1. Mechanism of action**

inhibits the enzyme dihydrofolate reductase, which in turn inhibits DNA synthesis

3.2. At a dose of 15 mg per week, taken orally or parenterally (subcutaneously, intramuscularly) is effective in terms of skin involvement (scleroderma), as well as in arthritis, myositis in the course of progressive systemic sclerosis.

4. Rituximab (Mabthera®)

4.1. Mechanism of action

– monoclonal anti-CD20 antibody, leads to depletion of B lymphocytes

4.2. At a dose of 375 mg/m² for four consecutive weeks in the first and sixth months has a good effect on skin tightness and pulmonary ventilatory parameters – forced vital capacity and diffusion capacity.

5. Cyclosporine

5.1. Inhibits T-cell activation

5.2. Proven efficacy in aggressive diffuse scleroderma

5.3. The most common side effect is nephrotoxicity.

6. Intravenous immunoglobulins (IVIg) – 2 gr/kg IVIg for six consecutive months

7. Plasmapheresis, photopheresis in diffuse cutaneous scleroderma

8. High doses of immunosuppressive drugs autologous stem cell transplantation.

9. Rapamycin – inhibits IL-2 dependent T-cell proliferation.

II. Antifibrotic drugs

1. Interferon α and Interferon γ

1.1. Interferon α – found to be ineffective in patients with early diffuse scleroderma (both cutaneous and pulmonary)

1.2. Interferon γ – moderate effect on skin manifestations (scleroderma) in patients with systemic sclerosis

2. D-penicillamine (Cuprenil®)

2.1. Mechanism of action

– antifibrotic effect, which is due to the fact that it blocks the binding of collagen fibers

– immunomodulatory effect?

2.2. Efficiency

– its effect has been demonstrated in a number of uncontrolled clinical trials in patients with early rapidly progressive scleroderma. It is established: 1. improvement in skin manifestations 2. reduction of mortality 3. reduction of the frequency of renal crises.

3. Recombinant human relaxin – reduces the synthesis of type I collagen by fibroblasts.

4. Minocycline.

5. Pirfenidone

5.1. Mechanism of action – blocks the activation of TGF- β as well as fibroblast growth factor in the lung.

5.2. Effective in idiopathic pulmonary fibrosis, not used in patients with scleroderma.

6. Tyrosine kinase inhibitors – Imatinib

6.1. Mechanism of action – inhibits TGF- β , leading to decreased synthesis of extracellular matrix proteins by fibroblasts.

6.2. In terms of efficacy, it was found to improve the modified Rodnan skin score as well as lung function.

7. Tocilizumab (RoActemra) – clinical trials in patients with early, diffuse scleroderma.

Table 10. Medicines used in patients with scleroderma and degree (level) of evidence

Medications	Degree (level) of proof
Immunomodulatory therapy	
Cyclophosphamide	A
Methotrexate	A
Mycophenolate	B
Rituximab	A, B
Immunoablation (high doses of Cyclophosphamide and subsequent stem cell transplantation)	A, C
Cyclosporine	C
Intravenous immunoglobulins	C
Photopheresis	A
Rapamycin	A
Antifibrotic therapy	
Interferon- α	A
Interferon- γ	C
D-penicillamine	A, B
Relaxin	A
Minocycline	B, C
Halofuginone	C
Pirfenidone	A
Imatinib	B

Level A: Randomized, placebo-controlled clinical trials; cohort studies; Level B: Retrospective, Exploratory cohort studies, extrapolation of level A results; Level C: Series of cases; extrapolation of level B results; Level D

III. Corticosteroids in progressive systemic sclerosis

The dose of corticosteroid administered to patients with scleroderma is determined by disease activity, the degree of skin involvement as assessed by the modified Rodnan skin score, visceral manifestations, and physician experience.

EULAR/EUSTAR recommendations for the treatment of various organ localizations in progressive systemic sclerosis. The recommendations are based on a combination of evidence established in randomized clinical trials and accepted consensus.

1. Phenomenon/Raynaud's syndrome

Raynaud's phenomenon (RP) is a clinical manifestation of vasospasm of the small arteries and arterioles, characterized by a three-phase color reaction of the fingers and toes, and sometimes of the ears or nose. Initially it is manifested by ischemia, then asphyxia and reactive hyperemia with changes in skin color from whitening to blue and finally redness in the restoration of blood flow caused by cold or emotional stress. The sharp demarcation between the affected and unaffected by a vasospasm area is typical.

Raynaud's phenomenon is classified as primary, in the absence of underlying disease or anatomical abnormality (Raynaud's disease). Secondary RP (Raynaud's syndrome) is observed in some rheumatic diseases: systemic lupus erythematosus (SLE), progressive systemic sclerosis (PSS), poly-

myositis/dermatomyositis (PM/DM), rheumatoid arthritis (RA), vasculitis, mixed connective tissue disease (MCTD).

Raynaud's phenomenon is found in 10% of the population, affecting 20% to 30% of young women, in the ratio of women: men – 9:1. The first symptoms of the disease usually appear during puberty, in young girls who have a familial predisposition.

1.1. Raynaud's primary phenomenon

The International Consensus on Diagnostic Criteria, Mavarakis et al., Is used to diagnose primary FR. since 2014.

Diagnostic criteria:

1. Vasospastic attacks caused by cold exposure or emotional stress.
2. Normal capillaroscopic picture.
3. Lack of digital ulcerations and gangrene, sclerodactyly, calcinosis, skin fibrosis.
4. Lack of systemic connective tissue disease.
5. Low titer negative or antinuclear antibodies (ANA);

Raynaud's disease is much more common in women than in men. The vasoconstrictive effects of estrogen may also be due to the tonic effect of the hormone on sympathetic nerve transmission, a direct effect on adrenergic signaling and/or an enhancement of the adrenergic response by vasoconstrictor prostanoids. The endothelium is actively involved in the regulation of vascular tone directly, processing signals from at least five different adrenoceptor subtypes (a2A/D, a2C, b1, b2, b3) or by releasing nitric oxide.

Table 11. Different dose regimens of corticosteroids and the most common side effects

Dosage regimen	The most common side effects
1. Pulse therapy with corticosteroid ≥ 250 mg daily for 1-5 days, usually 0.5-1 gr daily, administered intravenously for 1-3 days, 1 x monthly, after which therapy is continued with CS 30-60 mg daily	hypertension neuropsychiatric manifestations arrhythmias myalgias/artralgias
2. Very high dose corticosteroid ≥ 100 mg daily, administered intravenously, in divided doses, for no more than 1-2 weeks	psychosis risk of severe, complicated infections
3. High dose corticosteroid – 1 mg/kg for 2-4 weeks, then reduce the dose by 10% per week > 30 mg and ≤ 100 mg daily, administered intravenously or orally	signs of drug hypercorticism (Cushing's syndrome) - skin atrophy, acne hypertension hypokalaemia, hyperlipidemia, hyperglycaemia
4. Average dose of corticosteroid – 0.25-0.5 mg/kg daily for 2-4 weeks, administered intravenously > 7.5 mg and ≤ 30 mg daily, administered intravenously or orally	osteoporosis, osteonecrosis glaucoma, cataracts risk of infections insomnia, psychosis suppression of the hypothalamic pituitary-adrenal axisакцелерирана атеросклероза
5. Low dose corticosteroid ≤ 7.5 mg administered orally	osteoporosis, osteonecrosis suppression of the hypothalamic-pituitary-adrenal axis infections with a lower frequency compared to higher doses of CS
Alternating mode	reduction of side effects Osteoporosis occurs

Vasoconstriction induced by postsynaptic α_2 -adrenoceptors is strongly influenced by cold exposure in patients with Raynaud's phenomenon. An association between increased contractile response to α_2C agonists and increased protein tyrosine kinase activity and tyrosine phosphorylation has been established. Greater intracellular phosphorylation of tyrosine in response to cold exposure has been found in the arterioles of both primary and secondary RP patients. Elevated plasma levels of endothelin-1 are found in patients with MS compared to those with primary RP. In addition to its vasoconstrictor properties, endothelin-1 has been shown to promote fibrosis in patients with scleroderma. The levels of other substances produced by the endothelium have been shown to be abnormal in patients with secondary Raynaud's phenomenon. These include elevated levels of endothelin-1, angiotensin 2 (in diffuse scleroderma, but not limited), prothrombotic factor VIII/von Willebrand, as well as coagulation pathway activators and markers of impaired fibrinolysis. This may suggest that the endothelium plays an important role in the pathogenesis of Raynaud's secondary phenomenon by promoting vasoconstriction and thrombosis.

The central sympathetic system plays a role in the pathogenesis of RP because the local vibration applied to one arm causes vasoconstriction in the other, which is removed by proximal nerve block. In addition, vasoconstriction caused by emotional stimulus (stress) has been reported by many authors in patients with Raynaud's phenomenon, which suggests a component of the central nervous system.

1.2. The secondary Raynaud's phenomenon (Raynaud's syndrome) is characterized by a later age of onset, usually after 30 years, and patients are diagnosed with stigmas in the clinical picture, leading to systemic autoimmune disease, the presence of trophic changes in the fingers of the limbs, specific autoantibodies, pathological capillaroscopic finding. Among rheumatic diseases, the highest incidence is Raynaud's syndrome in progressive systemic sclerosis (90-95% of cases), usually preceding the onset of the disease by years, characterized by severe course and development of digital ulcers. The capillaroscopic finding in MS is characterized by the presence of dilated and giant capillaries, hemorrhages, avascular zones and neoangiogenesis.

M. Cutolo et al. describes three stages of capillaroscopic changes in SSc:

1. "Early" (Prescleroderma) – preserved position of the capillaries, without avascular areas, the appearance of several dilated and/or giant capillaries and several hemorrhages.

2. "Active" – changes in this stage include a larger number of giant capillaries and hemorrhages, a moderately reduced number of capillaries and in some cases diffuse pericapillary edema.

3. "Late" – it is characterized by extensive avascular areas, severe capillary rearrangement of the capillaroscopic field and branched capillaries.

When diagnosing Raynaud's phenomenon, it is extremely important to differentiate the primary from the secondary in rheumatic diseases. Additional tests are needed: capillaroscopy, acute phase parameters, ANA.

Table 12 presents medications used to treat Raynaud's syndrome and digital ulcers in patients with progressive systemic sclerosis.

A group of medications	Exemplary medications	Comments	Level of evidence
Medications for Raynaud's syndrome			
Calcium blockers	Nifedipine, Amlodipine, Diltiazem	Different response to different medications, slow dose titration reduces the frequency of side effects	A
ACE inhibitors	Quinipril	There is no evidence of efficacy	A
AT-1 blocker	Losartan	Effective in clinical trials, ineffective in clinical practice	A
5-HT antagonists	Ketanserine	Limited application	A
Selective serotonin reuptake inhibitors	Fluoxetine	Weaker vasodilatory effect compared to calcium channel blockers	A
Medications for digital ulcers			
Phosphodiesterase inhibitors	Sildenafil, Tadalafil	Improvement in the manifestations of Raynaud's syndrome and digital ulcerations	A
Parenteral vasodilators	Iloprost, Prostaglandin E, Treprostinil	Effective in digital ulcers	B, C
Endothelin receptor antagonists	Bosentan	Protects against the development of new ulcers	A
Statins		Reduce the frequency of digital ulcerations	A
Other remedies for digital ulcers		Local and systemic antibiotics, debridement, analgesics, use of healing techniques	
Surgical interventions - digital sympathectomy, sympathectomy with botulinum toxin (Botox), debridement.			

Level A: Randomized, placebo-controlled clinical trials; cohort studies; Level B: Retrospective, Exploratory cohort studies, extrapolation of level A results; Level C: Series of cases; extrapolation of level B results; Level D

I. EULAR/EUSTAR recommendations for scleroderma-associated vasculopathy.

1. Raynaud's syndrome

a) Dihydropyridine Ca²⁺ antagonist

• Nifedipine – as a drug of first choice; at a dose of 3 x 10-20 mg up to a maximum dose of 30-60 mg; reduces the frequency and severity of ischemic attacks.

b) Venous and oral prostanoids

• Iloprost – 0.5-2 ng/kg/min infusion in 3-5 consecutive days or 2 x 50-100 µg oral administration.

c) Peripheral vasodilators

d) Antiplatelet agents – Clopidogrel, Aspirin

Dihydropyridine Ca²⁺ antagonists are the drug of choice for scleroderma-related Raynaud's syndrome. In case of failure of their application, intravenous prostanoids are recommended.

2. Digital ulcerations

a) Iloprost – 0.5-2 ng/kg/min infusion on 3-5 consecutive days; accelerates healing of digital ulcers, as well as prevents the formation of new ulcers.

b) Bosentan – dual endothelin receptor antagonist; at a dose of 2 x 62.5 mg daily for 4 weeks, then 2 x 125 mg; prevents the development of new ulcerations.

Bosentan is recommended for diffuse systemic sclerosis with digital ulcerations, in the absence of effect from the use of prostanoids.

Recommendation	Strength of the recommendation
I. Scleroderma-associated digital vasculopathy	
Meta-analysis for the use of dihydropyridine Ca ²⁺ antagonists and a meta-analysis for the use of venous prostanoids. Nifedipine and intravenous Iloprost reduce the frequency and severity of ischemic attacks	A
Dihydropyridine Ca ²⁺ antagonists, usually Nifedipine, the drug of choice in SSR-Raynaud's syndrome, and intravenous prostanoids in severe Raynaud's syndrome, which developed in the course of progressive systemic sclerosis	
Two randomized clinical trials have shown that intravenous prostanoids, specifically intravenous Iloprost, are effective in healing digital ulcers in patients with SSc. Intravenous prostanoids should be used as a treatment for digital ulcers	A
Two randomized clinical trials show that Bosentan prevents the development of new digital ulcers, especially in patients with multiple ulcers on the fingers	A
Bosentan should be used in diffuse systemic sclerosis with multiple digital ulcerations following failure of Ca ²⁺ antagonists and prostanoids	

In Bulgaria according to the brief product characteristics:

I. Scleroderma-associated vasculopathy

1. Raynaud's syndrome

Dihydropyridines Ca²⁺ antagonists - not indicated for the treatment of scleroderma-related Raynaud's syndrome according to the brief characteristics of the product.

Venous prostanoids or parenteral vasodilators

Ilomedin 20 – concentrate for solution for infusion 1 ml/0.027 mg; 2 ml/0.067 mg. It is administered as an intravenous infusion at a dose of 0.5-2 ng/kg/min. According to the brief characteristics of the product, it is indicated for the treatment of severe Raynaud's disabling syndrome.

Peripheral vasodilators

• Naftidrofuril (Dusopharm) – 50 mg tablets; the recommended daily dose is 300 mg – 3 x 2 tablets; according to the brief characteristics of the product, it is indicated for the treatment of peripheral vascular disorders, including Raynaud's syndrome. The treatment can last from 1 to 6 months.

• Naftidrofuril (Dusodril) – 100 mg and 200 mg tablets. According to the brief characteristics of the product, it is indicated as a concomitant treatment for Raynaud's syndrome. The recommended daily dose is 2 x 200 mg daily for at least 3 months.

Antiplatelet agents – Aspirin, Clopidogrel.

Digital ulcerations

In the summary of product characteristics Ilomedin and Bosentan there is no therapeutic indication for digital ulcers in progressive systemic sclerosis.

In clinical practice in Bulgaria for digital ulcers in the course of progressive systemic sclerosis pulse therapy with Methylprednisolone 1000 mg and Cyclophosphamide 1000 mg intravenously for 6-12 months, as well as fractionated or unfractionated heparins.

Pulmonary arterial hypertension (PAH)

One of the serious complications of progressive systemic sclerosis is pulmonary arterial hypertension (PAH). PAH is more common in a limited form of progressive systemic sclerosis. Pulmonary arterial hypertension (PAH) is a condition that occurs with the presence of precapillary BH in the absence of other causes (such as BH as a result of lung disease).

Functional classification of PAH according to WHO:

Class I – without restriction of physical activity, normal physical activity does not cause fatigue, chest pain and syncope

Class II – slight limitation of physical activity, but without a problem at rest, normal physical activity causes fatigue, chest pain and discomfort

Class III – more pronounced restriction of physical activity, but without shortness of breath at rest, normal physical activity causes fatigue, chest pain and dyspnea

Class IV – pronounced signs of fatigue, dyspnea and chest pain during normal physical activity.

The incidence of PAH in patients with progressive systemic sclerosis is between 4.9-38%, on average 16%.

The clinical symptoms of PAH are represented by dyspnea on exertion (~ 60% of patients), fatigue or general weakness, general intolerance to physical exertion. As the disease worsens, dyspnoea at rest, chest pain (~ 40%), syncope (~ 40%) appear. In uncontrollable clinical manifestations of PAH, right heart failure develops, manifested by swelling of the lower extremities, enlarged jugular veins, hepatomegaly, progressive fatigue.

Recommendations for PAH screening (Sixth World PAH Symposium, 2017):

1. Annual PAH screening is recommended in asymptomatic patients with systemic sclerosis and other systemic connective tissue diseases.

2. Screening includes two steps - echocardiography and right cardiac catheterization for abnormalities in mean pulmonary arterial pressure.

3. The patient screening program should be part of a protocol or register.

Recommendations for the diagnosis of pulmonary arterial hypertension

1. Early diagnosis of PAH is performed through the following tests:

1.1. Lung radiography, functional examination of gas diffusion breathing, chest computed tomography to assess pulmonary fibrosis.

1.2. Measurement of the distance travelled in 6 minutes of walking on a flat surface (6 MWT)

1.3. Doppler echocardiography

Confirmation of the diagnosis is performed by right cardiac catheterization.

2. In the late stages of the disease are assessed:

Increasing severity of shortness of breath

Decreased tolerance of physical activity

Hoarseness

Hemoptysis

The goals of the treatment of PAH:

Indicators	Objectives of treatment
Functional class	I or II
Echocardiography /MRI of heart	Normal/close to normal right ventricular function
Hemodynamics	Normalization of the right ventricular function (right atrial pressure < 8 mm Hg and cardiac index > 2.5 to 3.0 l/min/m ²)
6MWT	> 380 to 440 m
Cardiopulmonary tests	Peak O ₂ consumption > 15 ml/min/kg and fan equivalent for CO ₂ < 45 l/min/l/min
Natriuretic peptide	Normal values

Recommendations for the treatment of pulmonary arterial hypertension

1. Symptomatic treatment

1.1. Diuretics – in manifestations of right heart failure

1.2. Continuous long-term oxygen supply

1.3. Oral anticoagulants

1.4. Digoxin

2. Specific therapy for PAH includes the following groups of drugs:

Calcium antagonists	Amlodipine, Diltiazem, Nifedipine
Prostanoids	Beraprost, Epoprostenol, Iloprost, Treprostinil,
Endothelin receptor antagonists	Ambrisentan, Bosentan, Macitentan
Phosphodiesterase type-5 inhibitors	Sildenafil, Tadalafil
Soluble guanylate cyclase stimulant	Riociguat

II. EULAR/EUSTAR recommendations for scleroderma-associated pulmonary arterial hypertension (SSc-PAH).

1. Endothelin receptor antagonists

Bosentan is a competitive endothelin receptor antagonist, with proven efficacy in pulmonary arterial hypertension developing in the course of progressive systemic sclerosis. The recommended daily dose is 2 x 62.5 mg for 4 weeks, followed by 2 x 125-250 mg. Improves physical capacity, functional class, some hemodynamic parameters and survival of patients.

Macitentan is a selective ETA receptor antagonist indicated for the treatment of PAH. Macitentan has been shown to improve long-term PAH in patients with systemic connective tissue disease.

2. Selective 5-phosphodiesterase inhibitors

Sildenafil is a selective phosphodiesterase inhibitor approved for the treatment of PAH II, III, IV functional class. A randomized clinical trial demonstrated its efficacy in SSc-PAH, showing improvement in physical capacity, functional class, and some haemodynamic parameters. The recommended dose is 3 x 20 mg, 40 mg, 80 mg, and is indicated in patients without sufficient effect of Bosentan. The mechanism of action is based on the selective inhibition of PDE5, which leads to smooth muscle relaxation, especially in the pulmonary vessels. This in turn causes vasodilation of the pulmonary vessels in patients with PAH, which improves mean pulmonary arterial pressure, pulmonary vascular resistance and cardiac index.

3. Intravenous Epoprostenol. FDA approved for the treatment of severe PAH (III, IV functional class), administered in a continuous infusion of 2 ng/kg/min.

It is contraindicated in severe left ventricular systolic dysfunction and pulmonary edema.

Intravenous iloprost – stable prostacyclin analogue, an alternative to the administration of intravenous epoprostenol, and in some countries it is preferred, despite the low class of evidence (C, IIa). The recommended dose is between 1 and 5 ng/kg/min, in isolated cases – up to 10 ng/kg/min.

Inhaler iloprost. It has been found that after a single inhaled dose of Iloprost, a reduction in mean pulmonary arterial pressure (ARI) of 10-20% was observed, lasting 45-60 minutes. The short duration of the effect requires frequent inhalations (6 to 12 times a day) to achieve a lasting effect with long-term use. Inhaled iloprost shows good tolerability. Treatment with inhaled Iloprost is indicated for the treatment of PAH III functional class according to NYHA. The dosage is between 6 times 2.5 µg and 9 times 5 µg/day.

Beraprost. This stable prostacyclin analogue for oral administration is approved for use in Japan and Korea in patients with primary and secondary PAH.

Treprostinil – it is administered subcutaneously or intravenously. Indicated for the treatment of patients with PAH with NYHA functional class II-IV.

4. **Riociguat** – soluble guanylate cyclase stimulant. It acts synergistically with nitric oxide, stimulates soluble guanylate cyclase, and the end result is vasodilation.

Combination treatment

Combination therapy is an option to counteract all the pathophysiological mechanisms that are present in PAH. It can be given by starting treatment with two (or more) medicines at the same time or by adding a second (or third) medicine to a previous therapy that has been found to be ineffective. It is currently unknown which of these two strategies is the better choice.

In many cases, only combined treatment with drugs from different groups leads to satisfactory clinical results. It can be expected that combination therapy, aimed at continuous improvement of PAH, will be the future standard and controlled studies have been launched for this purpose. Combination therapy, as a first step in the treatment regimen, is also recommended for decompensated right heart failure or threatened multiorgan failure (level of evidence C, level of recommendation IIa).

Algorithm of treatment from the V World Symposium on PAH 2013.

Функционален клас	FC II	FC III	FC IV
I (A or B)	Ambrisentan Bosentan Macitentan Riociguat Sildenafil Tadalafil	Ambrisentan, Bosentan, Epoprostenol i.v., Iloprost inhaled, Macitentan Riociguat, Sildenafil, Tadalafil, Treprostinil s.c., inhaled†	Epoprostenol i.v.
IIa (C)		Iloprost i.v.† Treprostinil i.v.	Ambrisentan, Bosentan, Iloprost inhaled and i.v.† Macitentan, Riociguat, Sildenafil, Tadalafil, Treprostinil s.c., i.v., inhaled†
IIb (B)		Beraprost†	
IIb (C)		Initial combination therapy	Initial combination therapy

The combined therapy for the treatment of pulmonary arterial hypertension can be presented as follows:

Recommendations	Strength of the recommendation
II. Scleroderma-associated pulmonary arterial hypertension (SSc-PAH)	
Two randomized clinical trials found that Bosentan improved physical capacity, functional class and some haemodynamic parameters. Indicated for the treatment of SSc-PAH	A/B
Two randomized clinical trials found that Sitaxentan improved physical capacity, functional class and some haemodynamic parameters. Also indicated for the treatment of SSc-PAH	A/B
A randomized clinical trial demonstrates that Sildenafil improves physical capacity, functional class, and some hemodynamic parameters in pulmonary arterial hypertension.	A/B
Sildenafil can be used to treat SSc-PAH	
A randomized clinical trial demonstrated that intravenous Epoprostenol improved physical capacity, functional class, and some hemodynamic parameters in pulmonary arterial hypertension. Indicated in severe SSc-PAH	A

In Bulgaria according to the brief product characteristics:

II. Scleroderma-associated pulmonary arterial hypertension

1. Endothelin receptor antagonists

Bosentan (Tracleer®) – tablets 62.5 mg and 125 mg, start 2 x 62.5 mg for 4 weeks, then increase the dose to 2 x 125 mg.

2. Selective 5-phosphodiesterase inhibitors

Sildenafil (Revatio®) – 20 mg tablets. The recommended daily dose is 3 x 20 mg.

III. EULAR/EUSTAR recommendations for scleroderma-associated skin involvement

Metotrexate. Two randomized clinical trials have demonstrated the effectiveness of MTX in terms of skin manifestations, assessed by the so-called. Rodnan skin score. Indicated for use in early diffuse scleroderma. The recommended dose is 15 mg/week.

Recommendations	Strength of the recommendation
III. Scleroderma-related skin involvement	
Two randomized clinical trials found that Metotrexate improved skin involvement in patients with early diffuse SSc. No positive effects on other organ damage have been demonstrated.	A
Metotrexate is recommended for the treatment of skin manifestations in patients with early diffuse systemic sclerosis	

IV. Biological therapy in progressive systemic sclerosis. Expects EMA approval in the next few years for a monoclonal antibody against IL-6 (Tocilizumab) in patients with progressive systemic sclerosis.

In Bulgaria according to the brief product characteristics:

III. Scleroderma-related skin involvement

1. In the brief description of **Metotrexate** there is no therapeutic indication for progressive systemic sclerosis. However, based on EULAR/EUSTAR recommendations, Metotrexate is used as an immunosuppressive therapy in patients with systemic sclerosis. In clinical practice, start with a dose of 10-15 mg/week, increasing the dose by 5 mg/week until a maximum dose of 25-30 mg/week is reached. The effect occurs between the 2nd and 3rd month. Metotrexate improves the skin score, assessing the degree of skin tightening, effective in arthritis, myositis, developing in the course of systemic sclerosis. The dosage forms available on the market are 2.5 mg tablets; 5 mg; 10 mg and vials for parenteral administration 50 mg/5 ml; 1000 mg/10 ml.

2. **D-penicillamine is an antifibrotic drug** with proven efficacy on the skin seal in scleroderma in a number of uncontrolled clinical trials. Systemic sclerosis is not included as a therapeutic indication in the summary of product characteristics. D-penicillamine is also not present in EULAR/EUSTAR recommendations for the treatment of progressive systemic sclerosis.

IV. EULAR/EUSTAR recommendations for scleroderma-related interstitial lung disease.

Cyclophosphamide according to two randomized clinical trials is recommended for the treatment of interstitial lung disease developing in the course of progressive systemic sclerosis. The recommended dose for oral administration is 1-2 mg/kg/day or for parenteral administration 600 mg/m² for 6 consecutive months. Treatment was then continued with another immunosuppressant Azathioprine 2.5 mg/kg. Cyclophosphamide increases total vital capacity, forced vital capacity, and improves quality of life in these patients. Does not affect the diffusion capacity for CO.

Recommendations	Strength of the recommendation
IV. Scleroderma-related interstitial lung disease	
Two randomized clinical trials have found that Cyclophosphamide should be used to treat interstitial lung disease that develops during systemic sclerosis, despite its toxicity.	A

In Bulgaria according to the brief product characteristics:

IV. Scleroderma-related interstitial lung disease

According to the summary of product characteristics Cyclophosphamide is indicated in patients with vasculitis and interstitial pulmonary fibrosis, fibrosing alveolitis developed in the course of systemic sclerosis. It is administered in a dose of 1-2 mg/kg/day. The dosage forms available on the market are 50 mg tablets; vials of 200 mg and 500 mg lyophilisate for injection.

* According to EULAR/EUSTAR recommendations, another immunosuppressive drug used in interstitial lung disease in scleroderma is Azathioprine. In our country, this therapeutic indication is not included in the brief description of Azathioprine.

V. EULAR/EUSTAR recommendations for scleroderma renal crisis

Recommendations	Strength of the recommendation
V. Scleroderma renal crisis	
There are no randomized clinical trials, experts recommend the use of ACE inhibitors to treat scleroderma renal crisis	C
Four retrospective studies concluded that steroids were associated with an increased risk of developing scleroderma renal crisis. Corticosteroid patients should be closely monitored for blood pressure and renal function.	C

VI. EULAR/EUSTAR recommendations for scleroderma-related gastrointestinal tract involvement

Recommendations	Strength of the recommendation
V. Scleroderma-related involvement of the gastrointestinal tract	
Despite the lack of randomized clinical trials, experts recommend PPIs (proton pump inhibitors) to prevent the development of gastroesophageal reflux disease, esophageal ulcers and structures	B
Despite the lack of randomized clinical trials, experts recommend prokinetics for the treatment of scleroderma-related motility disorders - dysphagia, gastroesophageal reflux, pseudoobstruction, etc.	C
Despite the lack of randomized clinical trials, experts recommend antibiotic therapy for malabsorption resulting from bacterial overgrowth.	D

One of the characteristic complications of progressive systemic sclerosis is the formation of soft tissue dystrophic calcifications.

One of the characteristic complications of progressive systemic sclerosis is the formation of soft tissue dystrophic calcifications.

Treatment of calcinosis in progressive systemic sclerosis	
Drug and dosage regimen	Probable mechanism
Warfarin 1 mg/day	reduces the levels of calcium-binding amino acids and carboxyglutamic acid reduces the uptake of technetium 99m-diphosphonate, found scintigraphically
Colchicine 1 mg/daily	anti-inflammatory effect
Probenecid 250-2000 mg/daily	increases renal phosphate clearance
Etidronate 10 mg/ daily	slows the progression of calcium deposits
Alendronate 10 mg/ daily	slows the progression of calcium deposits
Diltiazem 240 mg/ daily	Inhibits calcium influx in cells
Surgical excision	mechanical removal of calcifications
Aluminium hydroxide 1.68-2.24	reduces serum phosphate levels
Minocycline 50-100 mg/daily	inhibits matrix metalloproteinases, antibiotic effect, reduces calcium influx in cells
Topical application of a corticosteroid in calcium deposits	suppresses local inflammation
Анти-TNF inhibitors – 3-5 mg/kg	anti-inflammatory effect
Laser therapy with CO ₂	mechanical removal of calcifications

Eosinophilic fasciitis – a special form of progressive systemic sclerosis

It is a systemic disease that results from increased deposition of fibrous tissue in the dermis, hypodermis, subcutaneous adipose tissue, fascia and underlying muscles.

The disease occurs with involvement of the fascia of the limbs (fascia of the armpit, forearm, fascia lata), skin, internal organs. Clinically, the disease can occur with:

1. Fever

2. Pain – the pain of diffuse (eosinophilic) fasciitis is the result of the presence of an inflammatory infiltrate in the fascia, entheses, muscles, subcutaneous tissue and skin.

3. Arthralgia

4. Myalgias. Patients describe muscle pain throughout the body, but the focus is on the upper limbs and thighs along the fascia lata.

5. Arthritis.

6. Muscle contractures - they are common in the affected fascia.

7. Skin changes. Diffuse (eosinophilic) fasciitis is a disease that presents with scleroderma-like skin changes. The reason for this is the deposition of large amounts of extracellular matrix proteins in the dermis, hypodermis and subcutaneous adipose tissue. The fascia and regional muscles are additionally involved. The changes are localized mainly on the forearms, upper arms and lower limbs in the thighs.

8. Carpal tunnel syndrome. The syndrome is characteristic of chronic diffuse (eosinophilic) fasciitis.

itis. It is due to compression of the medial nerve in the wrist. The dystrophic changes typical of the syndrome gradually occur.

The disease occurs in three stages:

Clinical stage I – regional edema.

Clinical stage II – the skin of the limbs acquires the appearance of “orange peel” due to induration.

Clinical stage III – limited range of motion, flexion contractures and muscle atrophy.

The principles of treatment are confirmed by The European League Against Rheumatism.

Pharmacological therapy for diffuse (eosinophilic) fasciitis includes:

1. Corticosteroids

The recommended dose is 1 mg/kg for 1 month, followed by gradual dose reduction. The standard therapy lasted from 6 to 24 months.

2. Cytostatics and immunosuppressants – 5-Fluorouracil, Azathioprine (Imuran), Methotrexate, Cyclosporin A.

3. D-penicillamine.

4. Antimalarials

5. Mesenchymal inhibitors – Disulone

The prognosis for diffuse (eosinophilic) fasciitis is favorable. With strict adherence to the therapeutic regimen and established hygienic-dietary regime, long-term remission occurs.

References

- Amiri et al. Endothelium-restricted overexpression of human endothelin-1 causes vascular remodeling and endothelial dysfunction. *Circulation* 2004 12;110 (15):2233-40.
- Avouac J, Fransen J, Walker UA et al. Preliminary criteria for the very early diagnosis of systemic sclerosis: results of a Delphi Consensus Study from EULAR Scleroderma Trials and Research Group. *Ann Rheum Dis* 2011; 70: 476–481.
- Badesch DB et al. Pulmonary arterial hypertension: baseline characteristics from the REVEAL Registry. *Chest*. 2010 Feb;137 (2):376-87.
- Badesch DB, Abman SH, Ahearn GS, et al. Medical therapy for pulmonary arterial hypertension: ACCP evidence-based clinical practice guidelines. *Chest* 2004;126(1Suppl): 35S-62S.
- Badesch DB, Abman SH, Simonneau G, et al. Medical therapy for pulmonary arterial hypertension: updated ACCP evidence-based clinical practice guidelines. *Chest* 2007; 131:1917-28.
- Baines C, Pradeep K, Belch J. Raynaud phenomenon. *Rheumatology*; 2015.6; 1213-1218.
- Block JA, Sequeira W. Raynaud's phenomenon. *Lancet*; 2001; 357 (9273): 2042–2048.
- Bolster M, Siver R. Clinical features of systemic sclerosis. In *Rheumatology* edited by: Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. ELSEVIER, Toronto, Canada, 2011, 1373-1385.
- Campbell PM, EC LeRoy. Pathogenesis of systemic sclerosis: a vascular hypothesis. *Semin Arthritis Rheum* 1975; 4: 351-368.
- Clozel et al. Bosentan-improved cardiopulmonary vascular performance and increased plasma levels of endothelin-1 in porcine endotoxin shock. *BR J Pharmacol* 1996 Jun; 118 (3): 617–626.
- Conrad K, Schoesler W, Hiepe F, Fritzler MJ. eds. Autoantibodies in Systemic Autoimmune Diseases. A Diagnostic Reference. Pabst Science Publishers, 2007.
- Cutolo M, Sulli A, Pizzorni C et. al. Nailfold videocapillaroscopy assessment of microvascular damage in systemic sclerosis. *The journal of Rheumatology*. 2000; 27; 155-160.
- Denton CP. Pulmonary management of systemic sclerosis. In *Rheumatology* edited by: Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. ELSEVIER, Toronto, Canada, 2011, 1413-1425.
- Denton CP. Assessment of skin involvement in scleroderma. First EULAR/EUSTAR course.
- D'Angelo WA, Masi AT, Shulman LE. Pathologic observations in systemic sclerosis (scleroderma): a study of fifty-eight autopsy cases and fifty-eight matched controls. *Am J Med* 1969; 46: 428-440.
- Falanga V, Medsger TA Jr, Reichlin M, Rodnan GP. Linear scleroderma: clinical spectrum, prognosis and laboratory abnormalities. *Ann Intern Med* 1986; 104: 849-57.
- Fardoun MM, Nassif J, Issa K et al. Raynaud's Phenomenon: A Brief Review of the Underlying Mechanisms. *Front. Pharmacol.* 2016; 7:438.
- Fett N, Werth VP. Update on morphea: Part I. Epidemiology, clinical presentation, and pathogenesis. *J Am Acad Dermatol.* 2011; 64(2): 217–28. doi:10.1016/j.jaad.2010.05.045.
- Frank van den Hoogen et al. Classification Criteria for systemic sclerosis. An American College of Rheumatology/European League Against Rheumatism Collaborative Initiative. *Arthritis Rheum.*, 2013.
- Galiè et al. Treatment of patients with mildly symptomatic pulmonary arterial hypertension with bosentan (EARLY study): a double-blind, randomised controlled trial. *Lancet* 2008; 371, 2093-2100.
- Galiè N et al. Updated Treatment Algorithm of Pulmonary Arterial Hypertension *J Am Coll Cardiol* 2013; 62:D60-72.
- Goundry B, et al. Diagnosis and management of Raynaud's phenomenon. *BMJ*. 2012; 344:e289.
- Ho KT, Reveille JD. The clinical relevance of autoantibodies in scleroderma. *Arthritis Res Therapy* 2003; 5: 80-93.
- Hudson M, Taillefer S, Steele R et al. Improving the sensitivity of the American College of Rheumatology classification criteria for systemic sclerosis. *Clin Exp Rheumatol* 2007; 25: 754-7.
- Jing J, Dou TT, Yang JQ et al. Role of endothelin-1 in the skin fibrosis of systemic sclerosis. *European cytokine network*. 2015;26 (1); 10-14.
- Kreuter A, Krieg T, Worm M et al. German guidelines for the diagnosis and therapy of localized scleroderma. *J German Soc Dermatol JDDG*. 2016; 14(2): 199-216.

27. Kuhlmann et al. Endothelin-1-induced proliferation of human endothelial cells depends on activation of K⁺ channels and Ca²⁺ influx. *Acta Physiol Scand* 2005; Girgis et al; Decreased exhaled nitric oxide in pulmonary arterial hypertension: response to bosentan therapy. *Am J Respir Crit Care Med* 2005; 172 (3):352-7.
28. Lambova S. Raynaud's phenomenon. 2018. Chapter 3. <http://dx.doi.org/10.5772/intechopen.79362>
29. LeRoy EC, Medsger TA Jr. Criteria for the classification of early systemic sclerosis. *J Rheumatol*. 2001; 28: 1573-1576.
30. Lonzetti LS, Joyal F, Raynauld JP, et al. Updating the American College of Rheumatology preliminary classification criteria for systemic sclerosis: addition of severe nailfold capillaroscopy abnormalities markedly increases the sensitivity for limited scleroderma. *Arthritis Rheum*. 2001; 44: 735-736.
31. Matucci-Cerinic M, Allanore Y, Czirják L, et al. The challenge of early systemic sclerosis for the EULAR Scleroderma Trial and Research group (EUSTAR) community. It is time to cut the Gordian knot and develop a prevention or rescue strategy. *Ann Rheum Dis*. 2009; 68: 1377-1380.
32. Mayes M, Assassi S. Epidemiology and classification of scleroderma. In *Rheumatology* edited by: Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. ELSEVIER, Toronto, Canada, 2011, 1361-1371.
33. Maverakis E, Patel F, Kronenberg DG et al. International consensus criteria for the diagnosis of Raynaud's phenomenon. *J. Autoimmun.* 2014. 4, 60-65.
34. McGoon et al. Screening, early detection, and diagnosis of pulmonary arterial hypertension: ACCP evidence-based clinical practice guidelines. *Chest* 2004, 126(1 Suppl):14S-34S.
35. McLaughlin et al. Survival with first-line bosentan in patients with primary pulmonary hypertension. *Eur Respir J* 2005, 25, 1-6.
36. Nelson AM. Localized scleroderma including morphea, linear scleroderma, and eosinophilic fasciitis. *Curr Probl Pediatr*. 1996; 26 (9): 318-32.
37. Nihtyanova SI, Denton CP. Autoantibodies as predictive tools in systemic sclerosis. *Nat Rev rheumatol* 2010; 6: 112-116.
38. Rodnan GP, Benedek T. An historical account of the study of progressive systemic sclerosis (diffuse scleroderma). *Ann Intern Med* 1962; 567: 305-319.
39. Shoenfeld Y, Meroni PL. Eds. *The general Practice Guide to Autoimmune Diseases*, PABST Science Publishers, 2012.
40. Simonneau G, Galie N, Rubin LJ, et al. Clinical classification of pulmonary hypertension. *J Am Coll Cardiol* 2004; 43(12 Suppl S): 5S-12S.
41. Simms RW. Localized scleroderma and scleroderma-like syndromes. In *Rheumatology* edited by: Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. ELSEVIER, Toronto, Canada, 2011, 1433-1437.
42. Steen VD. The many faces of scleroderma. *Rheum Dis Clin North Am* 2008; 34: 1-15.
43. Steen VD. Management of systemic sclerosis. In *Rheumatology* edited by: Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. ELSEVIER, Toronto, Canada, 2011, 1403-1412.
44. Takehara K, Sato S. Localized scleroderma is an autoimmune disorder. *Rheumatology*, 2005; 44, 3,274-279.
45. Varga J, Lafyatis R. Pathogenesis of systemic sclerosis. In *Rheumatology* edited by: Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. ELSEVIER, Toronto, Canada, 2011, 1387-1402.
46. Wigley FM. Raynaud's phenomenon. *The New England Journal of Medicine*. 2002; 347(13):1001-1008.
47. Whitfield ML, Finley DR, Murray JI et al. Systemic and cell type-specific gene expression patterns in scleroderma skin. *Proc Natl Acad Sci USA* 2003; 100: 12319-12324.